

The value of the equation of state of dark energy

Raul Jimenez

*Department of Physics and Astronomy, University of Pennsylvania, Philadelphia,
PA 19104, USA*

Abstract

From recent CMB and Large Scale Structure observations the value of the equation of state of dark energy, assuming it to be constant in time, is constrained to be $-1.3 < w < -0.7$ at the 95% confidence level: consistent with dark energy being a classical vacuum term. Here we describe two novel and independent methods, sensitive to different systematics, that give the same value for w and similar confidence regions. This suggests that systematics are not an issue in current determinations of w . The first method yields a measurement of w that relies on the minimum number of model-dependent parameters; the second method is a non-parametric measurement of the time dependence of $w(z)$. We also present a method to statistically determine the edge of a distribution.

Key words:

1 Introduction

It is now established that the universe is accelerating (e.g., Dunlop et al. (1996); Spinrad et al. (1997); Riess et al. (1998); Perlmutter et al. (1999)): the next logical step is to unveil the nature of the accelerating force, for example, by determining its equation of state ($p = w(z)\rho$), where $w(z)$ need not to be constant and can vary with redshift.

During the past year we have seen a significant improvement in the accuracy of the equation of state, w measurements (assuming w to be constant in time, e.g. Spergel et al. (2003); Caldwell et al. (2003); Jimenez et al. (2003);

Email address: raulj@physics.upenn.edu (Raul Jimenez).

Tonry et al. (2003) and also Verde and Melchiorri articles in these proceedings). All these methods are sensitive to different systematics, yet the results are remarkably consistent: $-1.3 < w < -0.7$ at 95% confidence (Spergel et al., 2003). While this measurement will soon become even tighter with 2yr WMAP data and SDSS galaxy and Ly- α forest power spectrum release, it still assumes w to be constant in time. A significant challenge will be to accurately measure $w(z)$. There has been a number of papers discussing different methods to measure w (e.g. Alcock and Paczynski (1979); Turner and White (1997); Caldwell et al. (1998); Garavich et al. (1998); Birkinshaw (1999); Efstathiou (1999); Hui (1999); Newman and Davis (2000); Haiman et al. (2001); Huterer and Turner (2001); Wang and Garavich (2001); Weller and Albrecht (2001); Baccigalupi et al. (2002); Huterer (2002); Hu (2002b); Kujat et al. (2002); Hu (2002a); Lima and Alcaniz (2002)) while less attention has been devoted to the more challenging task of measuring $w(z)$ without parameterizations.

Below, we describe two novel methods: a) a measurement of w that relies on the minimum number of model dependent parameters. This method uses the determination of the position of the first acoustic peak in the CMB angular power spectrum (ℓ_1) and the absolute ages of galactic globular clusters (GCs). b) a non-parametric measurement of the time dependence of $w(z)$. This method is based on the *relative* ages of stellar populations. These techniques have been described in Jimenez and Loeb (2002); Jimenez et al. (2003)

2 w from the ℓ_1 -GCs ages method

If the universe is assumed to be flat, the position of the first acoustic peak (ℓ_1 in the standard spherical harmonics notation) depends primarily on the age of the universe and on the effective value¹ of w (Caldwell et al., 1998; Hu et al., 2001; Knox et al., 2001; Caldwell et al., 2003). As noted by these authors, for a fixed w value, a change in the physical density parameter $\omega_m = \Omega_m h^2$ that keeps the characteristic angular scale of the first acoustic peak fixed will also leave the age approximately unchanged. Thus an independent estimate of the absolute age of the universe at $z = 0$ combined with a measurement of ℓ_1 yields an estimate of w largely independent of other cosmological parameters (e.g., Ω_m and h).

This can be better understood by considering that for a constant $w(z)$, the

¹ I.e., the average value over redshift. We use $w(z)$ to indicate when w is allowed to vary over time.

age of a flat universe is given by

$$t_0 = H_0^{-1} \int_0^{\infty} (1+z)^{-1} \left[\Omega_{\text{rad}}(1+z)^4 + \Omega_m(1+z)^3 + \Omega_{\Lambda}(1+z)^{3(1+w)} \right]^{-1/2} dz. \quad (1)$$

The position of the first acoustic peak is fixed by the quantity

$$\theta_A = r_s(a_{\text{dec}})/d_A(a_{\text{dec}}), \quad (2)$$

where a_{dec} is the scale factor at decoupling, $r_s(a_{\text{dec}})$ is the sound horizon at decoupling, and $d_A(a_{\text{dec}})$ is the angular diameter distance at decoupling. For a flat universe,

$$r_s(a_{\text{dec}}) = \frac{c}{H_0 \sqrt{3}} \int_0^{a_{\text{dec}}} \left[\left(1 + \frac{3\Omega_b}{4\Omega_{\gamma}} \right) (\Omega_{\Lambda} x^{1-3w} + \Omega_m x + \Omega_{\text{rad}}) \right]^{-1/2} dx \quad (3)$$

and

$$d_A(a_{\text{dec}}) = \frac{c}{H_0} \int_{a_{\text{dec}}}^1 \left[\Omega_{\Lambda} x^{1-3w} + \Omega_m x + \Omega_{\text{rad}} \right]^{-1/2} dx. \quad (4)$$

See Verde et al. (2003) for more details. We use equations 1 to 4 and fix θ_A to be consistent with the *WMAP* determination ($\ell_1 \simeq 220$; Page et al., 2003). Figure 1 shows the allowed region in the age- w plane obtained for $\Omega_b h^2 = 0.02$, $0.05 < \Omega_m < 0.4$, and $0.5 < h < 0.9$. It is clear from the plot that an independent and accurate age determination of the universe will provide a measurement of w . Thus by using only the *WMAP* observation of ℓ_1 and an independent estimate of the age of the universe, one can place constraints on w , largely independent of other cosmological parameters. Note that this method is sensitive to a different redshift weighting than supernova and CMB measurements.

The ages of the oldest globular clusters (GCs) provide a lower limit to the total age of the universe. Since numerous star-forming galaxies have now been observed up to redshift $z = 6.6$ (e.g., Kodaira et al., 2003) and the oldest GCs contain the oldest stellar populations in galaxies, it is reasonable to assume that GCs too have formed at redshift $z > 6$. Krauss and Chaboyer (2003) perform the most careful analysis to date of the effects of systematics in GC age determinations. They estimate the age of the oldest Galactic GCs using the main-sequence turn-off luminosity and evaluate the errors with Monte-Carlo techniques, paying careful attention to uncertainties in the distance, to systematics, and to model uncertainties. They find a best-fit age for the




Fig. 1. The region of parameter space allowed if we fix θ_A (which in turns fixes the position of the first CMB acoustic peak) and $\omega_b \equiv \Omega_b h^2 = 0.02$, weakly constraining $0.05 < \Omega_m < 0.45$ and $0.5 < h < 0.9$.

oldest Galactic GCs of $12.5^{+3.4}_{-2.2}$ Gyr (95% confidence limits). We find that their probability distribution for the oldest GC age, can be accurately described by

$$P(t) = \frac{A}{\sigma(t-T)} \exp \left[-\frac{\ln [(t-T)/m]^2}{2\sigma^2} \right] \quad (5)$$

where t denotes the age of the oldest GCs, $A = 1.466$, $\sigma = 0.25$, $T = 6.5$ and $m = 5.9$. Since the oldest GCs likely formed at $z > 5$, for all reasonable cosmologies we only need add about 0.3 Gyr to the GC ages to obtain an estimate of the age of the universe. Only the oldest (low metallicity) GCs should be used in the age estimate, for which there is no age spread (see e.g. Rosenberg et al. (1999), which finds no age spread for GCs with metallicities $[Fe/H] < -1.2$).

In order to constrain w , we can now assume a flat universe and Monte Carlo simulate the distribution of ℓ_1 subject to only weak priors on the other cosmological parameters. For different values of w , Jimenez et al. (2003) generated 10^5 models, randomly sampling the cosmological parameters Ω_m , $\Omega_b h^2$ and h with uniform priors, $0.5 < h < 0.9$, $0.05 < \Omega_m < 0.45$, and $0.01 < \Omega_b h^2 < 0.03$. The next step is to impose an age of the universe constraint by randomly sampling these models so that the age of the universe has a probability distribution whose shape is given by equation 5, but offset by 0.3 Gyr. Jimenez et al. (2003) used the publicly available code **CMBFAST** (Seljak and Zaldarriaga, 1996) to compute the resulting distribution of ℓ_1 . This is shown in Figure 2 where the two solid lines are the 68% and 90% *joint* confidence levels.

As expected, the age alone does not constrain w ; there is a degeneracy between




Fig. 2. Confidence regions (68 and 90%) for the value of w are shown by the filled grey areas (marginalized over the position of the first acoustic peak in the CMB, ℓ_1). The solid lines denote contours which show the constraints in w using the derived age of the oldest Galactic globular clusters only. The dash and dotted lines are the the position of the first acoustic peak and confidence region as measured by *WMAP*. Note that the position of the first peak greatly constrains the value of w .

w and ℓ_1 . If we now use the *observed* position of the first acoustic peak as recently measured, in a model independent way, from *WMAP* (Page et al., 2003), we can constrain w with high accuracy. The filled contours in Figure 2 are marginalized values for w at the 68% and 90% confidence levels. Thus we find $w < -0.8$ ($w < -0.67$) at the 68% (90%) confidence level.

This determination depends solely on the GC determination of the age of the universe and on the *observed* position of the first acoustic peak in the CMB power spectrum. This constraint is slightly less stringent than that obtained by Spergel et al. (2003) from a joint likelihood analysis of *WMAP* with six external data sets (*WMAP* + CBI + ACBAR + 2dFGRS + Lyman α forest power spectrum + Type Ia supernovae + H_0 constraint from the *HST* key project), but is tighter than the CMB-only (*WMAP* + CBI + ACBAR) determination and comparable to the *WMAP* + ACBAR + CBI + *HST* constraint.

3 Measuring a variable equation of state for the dark energy

The popular approach for measuring $w(z)$ uses its effect on the luminosity distance of sources. In particular, the proposal for the Supernova/Acceleration Probe (SNAP) mission² suggests to monitor ~ 2000 Type Ia supernovae

² <http://snap.lbl.gov/>

across the sky per year and determine their luminosity distances up to a redshift $z \sim 1.5$ with high precision. However, the sensitivity of the luminosity distance to the redshift evolution of $w(z)$ is compromised by its integral nature (Maor et al. 2001),

$$d_L = (1+z) \int_z^0 (1+z') \frac{dt}{dz'} dz', \quad (6)$$

where $t(z)$ is the age of the Universe at a redshift z which depends on $w(z)$.

Jimenez and Loeb (2002) proposed an alternative method that offers a much better sensitivity to $w(z)$ since it measures the integrand of equation (6) directly. Any such method must rely on a clock that dates the variation in the age of the Universe with redshift. The clock is provided by spectroscopic dating of galaxy ages. Based on measurements of the age difference, Δt , between two passively-evolving galaxies that formed at the same time but are separated by a small redshift interval Δz , one can infer the derivative, (dz/dt) , from the ratio $(\Delta z/\Delta t)$. The statistical significance of the measurement can be improved by selecting fair samples of passively-evolving galaxies at the two redshifts and by comparing the upper cut-off in their age distributions. All selected galaxies need to have similar metallicities and low star formation rates (i.e. a red color), so that the average age of their stars would far exceed the age difference between the two galaxy samples, Δt .

This *differential age method* is much more reliable than a method based on an absolute age determination for galaxies (e.g., Dunlop et al. 1996; Alcaniz & Lima 2001; Stockton 2001). As demonstrated in the case of globular clusters, absolute stellar ages are more vulnerable to systematic uncertainties than relative ages (Stetson, Vandenberg & Bolte 1996). Moreover, absolute galaxy ages can only provide a lower limit to the age of the Universe and only place weak constraints on the possible histories of $w(z)$.

Consider a flat universe composed of matter and dark energy with an equation of state $p_Q = w(z)\rho_Q$. The Hubble parameter is $H^2 = H_0^2[\rho_T(z)/\rho_T(0)]$. Here, the subscripts Q , m , and T refer to the dark energy, the matter, or the total sum of the two, respectively. Assuming further that the matter is non-relativistic (i.e. effectively pressureless), we get

$$H_0^{-1} \frac{dz}{dt} = -(1+z)^{5/2} \left[\Omega_m(0) + \Omega_Q(0) \exp \left\{ 3 \int_0^z \frac{dz'}{(1+z')} w(z') \right\} \right]^{1/2}, \quad (7)$$

where we have used the energy conservation equation for the dark energy, $\dot{\rho}_Q = -3H(1+w(z))\rho_Q$ (Maor et al. 2000). Thus, (dz/dt) is related to the

equation of state of the dark energy through one integration only, while the luminosity distance in equation (6) is given by an integral of the inverse of equation (7), namely through two integrations. By differentiating (dz/dt) with respect to t we find,

$$H_0^{-2} \frac{d^2 z}{dt^2} = \frac{[H_0^{-1}(dz/dt)]^2}{(1+z)} \left[\frac{5}{2} + \frac{3}{2}w(z) \right] - \frac{3}{2}\Omega_m(0)(1+z)^4 w(z), \quad (8)$$

which depends *explicitly* on $w(z)$ without any integrations. Thus, the second derivative of redshift with respect to cosmic time measures $w(z)$ directly. While it is possible to find significantly different redshift histories of $w(z)$ for which the evolution of d_L is similar, this cannot be done for $(d^2 z/dt^2)$.

To obtain dz/dt , Jimenez and Loeb (2002) proposed to use the old envelope of the age–redshift relation of E/S0 galaxies. They assume that E/S0 at different redshifts are drawn from the same parent population with the bulk of their stellar populations formed at relatively high redshift (e.g., Bower et al., 1992; Stanford et al., 1998). Then, at relatively low redshift, they are evolving passively and may be used as “cosmic chronometers”.

Recently, Jimenez et al. (2003) applied this method using high-quality spectra of a sample of old stellar populations covering the largest possible range in redshift. To this aim they combined the following data sets: (i) the luminous red galaxy (LRG) sample from the Sloan Digital Sky Survey (SDSS) early data release (Eisenstein et al., 2001); (ii) the sample of field early-type galaxies from Treu et al. (1999, 2001, 2002, hereafter the Treu et al. sample); (iii) a sample of red galaxies in the galaxy cluster MS1054–0321 at $z = 0.83$; and (iv) the two radio galaxies 53W091 and 53W069 (Dunlop et al., 1996; Spinrad et al., 1997; Nolan et al., 2003, Dey et al., in preparation). They obtained the age of the dominant stellar population in the galaxies by fitting single stellar population models (Jimenez et al., 1998) to the observed spectrum.

Figure 3 shows the derived single-burst equivalent ages of the galaxies as a function of redshift. The circles correspond to galaxies in the SDSS LRG sample, triangles refer to the Treu et al. sample, diamonds are galaxies in MS1054–0321, and crosses are 53W091 and 53W069. For clarity, the SDSS LRG points have not been plotted for ages < 7 Gyr. Typical errors on the ages of LRG galaxies are 10% and are not plotted. An age–redshift relation (i.e., an “edge” or “envelope” of the galaxy distribution in the age–redshift plane) is apparent from $z = 0$ to $z = 1.5$. Jimenez et al. (2003) also show that small recent episodes of star formation do not significantly bias the age determination.

What are the observational limits that can be imposed on $w(z)$? As a consistency check we note that the age at $z = 0$ obtained with this method is




Fig. 3. Top panel: Age-redshift envelope obtained for the galaxies studied in Jimenez et al. (2003). Open circles correspond to galaxies from the SDSS LRG sample, triangles to the Treu et al. sample, diamonds to MS1054–0321, and crosses are 53W091 and 53W069. A clear trend is present: galaxies age as the redshift decreases. The overall shape of this trend is in fair agreement with theoretical expectations for a Λ CDM (solid line) with $\Omega_m = 0.27$ and $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The dotted line represent an alternative model disfavored by the data: $w(z) = -2$ for $z > 1$ and then grows linearly from $w = -2$ at $z = 1$ to $w = 0$ at $z = 0$. Bottom panel: histogram along the age axis of top panel for the redshift range $0 < z < 0.04$ and $0.08 < z < 0.12$. The clear shift between the two histograms is a measurement of dz/dt and therefore allows us to measure H_0 .

in good agreement with GC ages and that the Treu et al. galaxy ages agree with those from of SDSS LRG sample where they overlap. The solid line in Figure 3 corresponds to the age-redshift relation for a flat, reference Λ CDM model: $\Omega_m = 0.27$, $h = 71$, and $w = -1$ (WMAP best fit model). This is consistent with the observed age-redshift envelope, and it seems to indicate that we live in a universe with a classical vacuum energy density and that, on average, stars in the oldest galaxies formed about 0.7 Gyr after the Big Bang (e.g., at redshifts $z \sim 7 - 10$). To make the observed age-redshift relation consistent with a model with widely different $w(z)$ behavior, one would have to infer that early-type galaxies at different redshifts have very different

formation epochs for their stellar populations. For example, the dotted line in Figure 3 indicates the age-redshift relation for a passively evolving population in a universe where $w(z) = -2$ for $z > 1$ and grows linearly from $w = -2$ at $z = 1$ to $w = 0$ at $z = 0$. For this model to work, we would have to conclude that the oldest galaxies at $z \sim 1.5$ formed about 0.7 Gyr after the Big Bang, but that the oldest galaxies at $z \sim 0$ formed ~ 3 Gyr after the Big Bang. Since the oldest galaxies in the LRG sample are as old as the oldest GCs, we consider this to be an unlikely explanation. Furthermore, not only would this scenario require that *all* local galaxies either formed their stars more recently than galaxies at high redshift or had recent episodes of star formation, but would also require a remarkable fine-tuning for them all to end up with exactly the same age. We therefore conclude that this extreme model for $w(z)$ is unlikely given the data.

Unfortunately, the small number of galaxies in Jimenez et al. (2003) samples at $z > 0.2$ did not allow them to compute dz/dt with enough accuracy to constrain $w(z)$, and therefore this measurement will have to await better data. However, the $z < 0.2$ region is well populated by LRG galaxies (Figure 3) and they determined dt/dz at $z \sim 0$. This is illustrated in the bottom panel of Figure 3 by the clear shift in the upper envelope of the age histogram between the two redshift ranges $0 < z < 0.04$ and $0.08 < z < 0.12$. This can be used to determine H_0 , which relies upon determining the “edge” of the galaxy age distribution in different redshift intervals.

If the ages of the galaxies were known with infinite accuracy, for each galaxy i at redshift z_i , one could associate an age a_i . The probability for the age of the oldest stellar population in galaxy i , $P(t|a_i)$, would be given by a step function which jumps from 0 to 1 at a_i . An age-redshift relation edge could then be obtained by dividing the galaxy sample into suitably-large redshift bins and multiplying the $P(t|a_i)$ for all the galaxies in each bin.

In practice, the age of each galaxy is measured with some error δa_i . We thus assume $P(t|a_i) = 1$ if $t > a_i + \delta a_i$ and $\ln P(a_i) = -x^2$ where $x = (a_i + \delta a_i - t)/(\sqrt{2}\delta a_i)$ otherwise. Jimenez et al. (2003) divided the $z < 0.2$ portion of the LRG sample into 51 redshift bins. For each bin, they obtained $P(z, t)$ by multiplying the $P(t|a_i)$ of the galaxies in that bin. They define the “edge” of the distribution $t(z)$ to be where $\ln P(z, t)$ drops by 0.5 from its maximum (i.e. $\Delta \ln P = 0.5$ and associate an error to this determination $\delta_{t(z)}$ given by $t_2(z) - t(z)$ where $t_2(z)$ corresponds to where $\ln P(z, t)$ drops by 2 from its maximum (i.e. $\Delta \ln P = 2$, this approximately corresponds to the 68% confidence level). This procedure makes the determination of the “edge” less sensitive to the outliers (see also Raychaudhury et al. (1997)). Figure 4 presents the resulting $t(z)$ relation.




Fig. 4. The binned, low-redshift age-redshift relation, derived from the SDSS LRG sample. The solid line is a best-fit to the edge.




Fig. 5. The value for H_0 derived from Figure 4 as a function of redshift cut-off for computing the slope. Note that it is very weakly dependent on the value of this cut-off. The solid circle is the adopted value. The shaded are corresponds to the 1σ confidence region from the Hubble Key Project. For $z > 0.18$ the probability of the fit being a straight line is $P_{\geq\chi^2} < 0.1$, while at $z = 0.17$ is 0.32.

For $z < z_{\max}$, Jimenez et al. (2003) fit $t(z)$ with a straight line whose slope dt/dz is related to the Hubble constant at an effective redshift by $H(z_{\text{eff}}) = -\frac{1}{1+z_{\text{eff}}}(\frac{dt}{dz})^{-1}$. This fit is performed by standard χ^2 minimization. They also compute $P_{\geq\chi^2}$, the probability of obtaining equal or greater value of the reduced χ^2 if the $t(z)$ points were truly lying on a straight line. Values of $P_{\geq\chi^2} < 0.1$ means that a straight line is not a good fit to the points.

For $w = -1$, $H_0 = H(z_{\text{eff}}) [\Omega_m(1 + z_{\text{eff}})^3 + \Omega_\Lambda]^{-\frac{1}{2}}$ where $\Omega_m = 0.27$, $\Omega_\Lambda = 0.7$. Figure 5 shows how the H_0 measurement depends on z_{max} . For $z_{\text{max}} > 0.17$ not only the value for H_0 drifts but also a straight line is not a good fit to the points ($P_{\geq \chi^2}$ suddenly drops below 0.1). For their H_0 determination, $z_{\text{max}} = 0.17$, $P_{\geq \chi^2} = 0.32$, $z_{\text{eff}} = 0.09$, and the correction from $H(z_{\text{eff}})$ to H_0 is a 4% effect. They obtained $H_0 = 69 \pm 12 \text{ km s}^{-1} \text{ Mpc}^{-1}$. Which is in good agreement with the Hubble Key Project measurement ($h = 0.72 \pm 0.03 \pm 0.07$; Freedman et al., 2001), with the valued derived from the joint likelihood analysis of *WMAP* + 2dFGRS + Lyman- α forest power spectrum ($h = 0.71_{-0.08}^{+0.04}$; Spergel et al., 2003), with gravitational lens time delay determinations ($h = 0.59_{-0.07}^{+0.12} \pm 0.03$; Treu and Koopmans, 2002), and Sunyaev-Zeldovich measurements ($h = 0.60_{-0.04}^{+0.04+0.13}$; Reese et al., 2002).

This check at $z = 0$ gives some confidence about the possibility of measuring $w(z)$ when larger samples of elliptical galaxies at higher redshifts become available in the near future. DEEP2 and the full SDSS catalog will provide excellent datasets for doing this.

I warmly thank my collaborators Avi Loeb, Dan Stern, Tommaso Treu and Licia Verde.

References

- Alcock, C., Paczynski, B., 1979. *Nature* 281, 358.
- Baccigalupi, C., Balbi, A., Matarrese, S., Perrotta, F., Vittorio, N., 2002. *PRD* 65, 63520.
- Birkinshaw, M., 1999. The Sunyaev-Zel'dovich Effect: an Update. In: *AIP Conf. Proc.* 476: 3K cosmology. pp. 298.
- Bower, R. G., Lucey, J. R., Ellis, R. S., 1992. *MNRAS* 254, 601.
- Caldwell, R., Kamionkowski, M., Weinberg, N., 2003. *astro-ph/0305008*.
- Caldwell, R. R., Dave, R., Steinhardt, P. J., 1998. *PRD* 80, 1582–1585.
- Dunlop, J., Peacock, J., Spinrad, H., Dey, A., Jimenez, R., Stern, D., Windhorst, R., 1996. *Nature* 381, 581–584.
- Efstathiou, G., 1999. *MNRAS* 310, 842–850.
- Eisenstein et al., D. J., 2001. *AJ* 122, 2267–2280.
- Freedman, W. L. et al., 2001. *ApJ* 553, 47–72.
- Garnavich et al., P. M., 1998. *ApJ* 509, 74.
- Haiman, Z., Mohr, J. J., Holder, G. P., 2001. *ApJ* 553, 545–561.
- Hu, W., 2002a. *PRD* 66, 83515.
- Hu, W., 2002b. *PRD* 65, 23003.
- Hu, W., Fukugita, M., Zaldarriaga, M., Tegmark, M., 2001. *ApJ* 549, 669–680.
- Hui, L., 1999. *ApJL* 519, L9–L12.
- Huterer, D., 2002. *PRD* 65, 63001.

- Huterer, D., Turner, M. S., 2001. PRD 64, 123527.
- Jimenez, R., Loeb, A., 2002. ApJ 573, 37–42.
- Jimenez, R., Padoan, P., Matteucci, F., Heavens, A. F., 1998. MNRAS 299, 123.
- Jimenez, R., Verde, L., Treu, T., Stern, D., 2003. astro-ph/0302560.
- Knox, L., Christensen, N., Skordis, C., 2001. ApJL 563, L95–L98.
- Kodaira et al., K., 2003. astro-ph/0301096.
- Krauss, L. M., Chaboyer, B., 2003. Science 299, 65–70.
- Kujat, J., Linn, A. M., Scherrer, R. J., Weinberg, D. H., 2002. ApJ 572, 1–14.
- Lima, J. A. S., Alcaniz, J. S., 2002. ApJ 566, 15–18.
- Newman, J. A., Davis, M., 2000. ApJL 534, L11–L14.
- Nolan, L., Dunlop, J., Jimenez, R., Heavens, A. F., 2003. MNRAS 341, 464.
- Page et al., L., 2003. astro-ph/0302220.
- Perlmutter et al., S., 1999. ApJ 517, 565–586.
- Raychaudhury, S., von Braun, K., Bernstein, G. M., Guhathakurta, P., 1997. AJ 113, 2046–2053.
- Reese, E. D., Carlstrom, J. E., Joy, M., Mohr, J. J., Grego, L., Holzappel, W. L., 2002. ApJ 581, 53.
- Riess et al., A. G., 1998. AJ 116, 1009–1038.
- Rosenberg, A., Saviane, I., Piotto, G., Aparicio, A., 1999. AJ 118, 2306–2320.
- Seljak, U., Zaldarriaga, M., 1996. ApJ 469, 437.
- Spergel et al., D., 2003. astro-ph/0302209.
- Spinrad, H., Dey, A., Stern, D., Dunlop, J., Peacock, J., Jimenez, R., Windhorst, 1997. ApJ 484, 581.
- Stanford, S. A., Eisenhardt, P. R. M., Dickinson, M., 1998. ApJ 492, 461.
- Tonry et al., J., 2003. astro-ph/0305008.
- Treu, T., Koopmans, L. V. E., 2002. MNRAS 337, L6–L10.
- Treu, T., Stiavelli, M., Casertano, S., Møller, P., Bertin, G., 1999. MNRAS 308, 1037–1052.
- Treu, T., Stiavelli, M., Casertano, S., Møller, P., Bertin, G., 2002. ApJL 564, L13–L16.
- Treu, T., Stiavelli, M., Møller, P., Casertano, S., Bertin, G., 2001. MNRAS 326, 221.
- Turner, M. S., White, M., 1997. PRD 56, 4439.
- Verde et al., L., 2003. astro-ph/0302218.
- Wang, Y., Garnavich, P. M., 2001. ApJ 552, 445–451.
- Weller, J., Albrecht, A., 2001. PRL 86, 1939–1942.